(Contents continued from back cover)

The transit rate of different-sized model dosage forms through the human colon and the effects of a lactulose-induced catharsis
P.J. Watts, L. Barrow, K.P. Steed, C.G. Wilson, R.C. Spiller, C.D. Melia and M.C. Davies (UK) 215
Solid-phase extraction of naftidrofuryl from human plasma for high-performance liquid chromatography analysis
P.J. Waaler and B.W. Müller (Germany, Norway) 223
Biopharmaceutical studies of naftidrofuryl in hydrocolloid matrix tablets
P.J. Waaler, C. Graffner and B.W. Müller (Germany, Norway, Sweden) 229
Aqueous nanodispersions prepared by a salting-out process
H. Ibrahim, C. Bindschaedler, E. Doelker, P. Buri and R. Gurny (Switzerland) 239
Preparation of aqueous polymeric nanodispersions by a reversible salting-out process: influence of process parameters on particle size
E. Allémann, R. Gurny and E. Doelker (Switzerland) 247
Release of cromoglycate from dextran and inulin conjugates
A.S. Williams and G. Taylor (UK) 255
Surfactant effects in percutaneous absorption. I. Effects on the transdermal flux of methyl nicotinate
P. Ashton, K.A. Walters, K.R. Brain and J. Hadgraft (USA, UK) 261
Surfactant effects in percutaneous absorption. II. Effects on protein and lipid structure of the stratum corneum
P. Ashton, K.A. Walters, K.R. Brain and J. Hadgraft (USA, UK) 265

Author Index 271

Noticeboard 275